

INFESTATION OF WORMS IN ADULT PATIENTS WITH ANAEMIA

Lubna Adnan, Rizwan Ali Talpur,
Rabnawaz Sathio, Muhammad Wasif Saleem

ABSTRACT

OBJECTIVE: Main objective of this study was to see infestation of worms in adult patients with anaemia.

METHODOLOGY: This was cross-sectional, descriptive study conducted at Department of Pathology, Chandka Medical College, Larkana for the period of 12 months (November 2019 to November 2020). Stool analysis for anaemic patients and controls was performed to detect infestation of worms. Data was analyzed using SPSS 24.0

RESULTS: A total of 400 patients were selected for determining the worm infestation frequency in adult patients with anemia. Out of 400 patients, 179 (44.75%) were aged between 18-30 years, 128 (32%) were between 31-40 years, and 93 (23.25%) were between 40-50 years. The mean age of patients was 28.44 ± 3.23 years. Male to female ratio was 1:2. The worm infestation frequency was 46.25%. Anemia was seen in 47 patients

CONCLUSION: This study explored good frequency rate of worm infestation in association with anaemia. Our results were mostly in supported by results of other studies. Higher authorities should take more serious steps in prevention of such parasitic diseases by conducting surveys, and to control the spread of parasites.

Keywords: Worms infestation, anaemia, adult population, stool examination.

How to cite this: Saleem W M, Shaikh H, Pirzado A A, Mahar S, Association Of Vitamin D Deficiency With Hypothyroidism. JIMC 2021; 4.(2) :201-203

Correspondence:

Dr. Lubna Adnan,
Assistant Professor
Department of Pathology
Chandka Medical College SMBBMU Larkana
Email: roma.adnan@gmail.com

INTRODUCTION

Anaemia is described as decreased levels of haemoglobin, caused by acquired or genetic changes in number or shape of the red blood cells. (1) Many causes are observed which cause anaemia including nutritional deficiencies, gastrointestinal bleed, genitourinary bleed, malabsorption syndromes, medication, increased destruction of blood due to acquired or genetic causes and inflammatory diseases. (2)

In developing countries, infestation by parasitic worms or Helminths is common aetiology. They cause anaemia by gastrointestinal bleeding or cause decreased micronutrient absorption. (3-4) Helminths which are transmitted from the soil have been found to cause between 5-39 million morbid lifestyles, leading to reduced psycho-physical growth and anaemia. (5) Infestation by worms has been associated with poor quality of life; therefore, it is more prevalent in developing and under-developed countries. (6)

The prevalence of infections by intestinal Helminths is observed low (e.g., 5.3%) as compared to protozoan infections (e.g., 16.7 – 18%) globally. (7) More than 1 billion individuals residing in under-developed countries or in areas with no or less availability of clean and sterile water are infected with intestinal Helminths. (8) The ratio is also high in counties with over-crowding e.g., China, India, and Pakistan etc. Globally more

than one billion populations are affected with Ascariasis and in India alone; more than 100 million individuals are infected. It is also postulated that over 20 million individuals are infected with Ascariasis in Pakistan. (9)

Additionally, *Trichuris trichiura* (whipworm), *Necator americanus* (hookworm) and *Ancylostoma duodenale* (hookworm) are also common. (10) Guidelines regarding these worm infestations and their management had been published by World Health Organization (WHO) in 2011. They recommended the use of anti-helminthic drugs (e.g., albendazole or mebendazole) should be used for the management and prevention of worm infestations in endemic areas. (11) Association of anaemia with worm infestations is studied limitedly; therefore, we designed this study to observe the presence of anaemia in patients with worm infestations in adult population.

PATIENTS AND METHODS

This was a cross-sectional, descriptive study conducted at Department of Pathology, Chandka Medical College, Larkana. The study was carried out for the period of 12 months (November 2019 to November 2020). 400 stool examinations were performed. A total of 185 patients were selected for the study. Patients aged between 18 to 50 years, both males and females were included using non-probability consecutive sampling. Patients with bleeding disorders, haemolytic anaemia, aplastic anaemia, leukaemia/lymphomas were excluded from the study. Cut-off value for anaemia was 13.5 g/dL for males and 11.5 g/dL for females. Patients

were labelled as worm infested if stool examination was positive for any cyst, ova eggs and/or worms. After getting approval from Ethical Committee, the process was explained, and informed consent was taken from all patients. The stool examination was performed by Pathologist in the laboratory. All data was recorded and analysed using SPSS 24.0. Means and standard deviations were calculated for quantitative variables e.g., haemoglobin levels, age, and frequencies etc. Chi-square test was observed to see the correlation of anaemia with worm infestation.

Stool Examination	Number	Percentage
Cysts	65	35.71%
Ova	39	21.08%
Eggs	21	11.35%
Larvae	42	22.70%
Worms	18	9.72%
Ancylostoma duodenale	3	16.66%
Ascaris lumbricoides	11	61.11%
Trichuris trichura	3	16.66%
Taenia saginata	1	5.55%

Worm Infestations	18-30 Years	31-40 Years	40-50 Years	Total
Yes	87	62	36	185
No	92	66	57	215

Stool Examination	Anaemia	
	Present	Absent
Cysts	18	47
Ova	5	34
Eggs	2	19
Larvae	11	33
Worms	11	7

DISCUSSION

Infestation of worms is major pandemic global health issue, which affect a large population. Additionally, as compared to developed countries, it is more common in developing countries. ⁽¹²⁻¹³⁾Among all parasitic infections, Ascaris has more importance because of its high prevalence rate. As per estimation, it is considered that more than one billion individuals are affected by this parasite globally. Likewise, the prevalence of Ascaris is over 140 million in India, over 86 million in China and over 21 million in Pakistan, making it very

RESULTS

A total of 400 patients were selected for determining the worm infestation frequency in adult patients with anaemia. Out of 400 patients, 179 (44.75%) were aged between 18-30 years, 128 (32%) were between 31-40 years, and 93 (23.25%) were between 40-50 years. The mean age of patients was 28.44 ± 3.23 years. Male to female ratio was 1:2. The worm infestation frequency was 46.25%. Anaemia was seen in 47 patients (Tables 1 essential health concern in these countries. ⁽¹⁴⁾Due to these problems, this study was conducted to see its association with anaemia. The worm infestation frequency in our study was 46.25%. Additionally, the infestation by various worms included Ascaris lumbricoides (61.66%), Ancylostoma duodenale

CONCLUSION

This study explored good frequency rate of worm infestation in association with anaemia. Our results were mostly in supported by results of other studies. Higher authorities should take more serious steps in prevention of such parasitic diseases by conducting surveys, and to control the spread of parasites.

Conflict of interest: None.

Funding Source: None.

REFERENCES

- Gardner W, Kassebaum N. Global, Regional, and National Prevalence of Anemia and Its Causes in 204 Countries and Territories, 1990-2019. *Current Development in Nutrition*. 2020;4(Suppl 2):830.
- Endris BS, Dinant G, Gebreyesus SH, Spigt M. Risk factors of anemia among preschool children in Ethiopia: a Bayesian geo-statistical model. *BMC Nutr*. 2022;8:2.
- Banda GT, Deribe L, Davey G. How can we better integrate the prevention, treatment, control and elimination of neglected tropical diseases with other healthy interventions? A systemic review. *BMJ Global Health*. 2021;6:e006968.
- Mengist HM, Zewdie O, Belew A. Intestinal helminthic infection and anemia among pregnant women attending ante-natal care (ANC) in East Wollega, Oromia, Ethiopia. *BMC Res Notes*. 2017;10:440.
- Nery SV, Pickering AJ, Abate E, Asmare A, Barrett L, Benjamin-Chung J et al. The role of water, sanitation and hygiene interventions in reducing soil-transmitted helminths: interpreting the evidence and identifying next steps. *Parasites Vectors*. 2019;12:273.
- Kwong LH, Sen D, Islam S, Shahriar S, Benjamin-Chung J, Arnold BF et al. Effect of sanitation improvements on soil-transmitted helminth eggs in courtyard soil from rural Bangladesh: Evidence from a cluster-randomized controlled trial. *PLoS Negl Trop Dis*. 2021;15(7):e0008815.
- Punsawad C, Phasuk N, Bunratsami S, Thongtup K, Siripakonuaong N, Nongnau S. Prevalence of intestinal parasitic infection and associated risk factors among village health volunteers in rural communities of southern Thailand. *BMC Public Health*. 2017;17:564.
- Lo AC, Faye B, Gyan BA, Amoah LE. Plasmodium and intestinal parasite perturbations of the infected host's inflammatory responses: a systemic review. *Parasitic Vectors*. 2018;11:387.
- Ali SA, Niaz S, Aguilar-Marcelino L, Ali W, Ali M, Khan A et al. Prevalence of Ascaris lumbricoides in contaminated faecal samples of children residing in urban areas of Lahore, Pakistan. *Sci Rep*. 2020;10:21815.
- Zelege AJ, Bayih AG, Afework S, Gilleard JS. Treatment efficacy and re-infection rates of soil-transmitted helminths following mebendazole treatment in schoolchildren, Northwest Ethiopia. *Trop Med Health*. 2020;48:90.
- Gandasegui J, Martinez-Valladares M, Grau-Pujol B, Krolewiecki AJ, Balaria-Fouce R, Gelaye W et al. Role of DNA-detection-based tools for monitoring the soil-transmitted helminth treatment response in drug-efficacy trials. *PLoS Negl Trop Dis*. 2020;14(2):e0007931.

12. Mona AK. Frequency of intestinal parasitic infestation in children of 5-12 years of age in Abbottabad. J Ayub Med Coll Abbottabad 2003;15(2):28-30.

13. Wani SA, Ahmed F, Zargar SA, Dar PA, Dar ZA, Jan TR. Intestinal helminthes in a population of children from Kashmir valley, India. J Helminthol 2008;82:313-7.

14. Alam M, Khattak AI, Talha M. Anemia and intestinal parasitic infestation in school children in Skardu. Pak Armed Forces Med J 2007;57(1):77-81.

15. Shad JA, Lee YR. Pancreatitis due to Ascarislumbricoides: Second occurrence after 2 years. South Med J 2001;94:78-80.

16. Khan A, Sultan A, Dan AMK, Rashid H, Najmi SAA. A study of prevalence, distribution and risk factors of intestinal helminth infestation in District Bagh (Azad Kashmir). Pak Armed Forces Med J 2004;54:243-8.

17. Khan W, Nisa NU, Khan A, Naqvi SMHM. Endemicity of intestinal parasites with special reference to nematodes in individuals related to education (students, staff & workers) in Swat KP, Pakistan. Pak J Nematol 2012;30(1):77- 85.

Authors

1. Dr .Rizwan Ali Talpur
Mphil Pathology
Assistant Professor
Department of Pathology, Chandka Medical College
SMBBMU Larkana
2. Dr. Rabnawaz Sathio
Mphil Physiology
Assistant Professor
Department of Physiology, Roshan Suleman Medical
College, Tando Adam
3. Dr. Muhammad Wasif Saleem
Mphil Pathology
Assistant Professor
Department of Pathology, Liaquat University of Medical and
Health Sciences, Jamshoro.

<i>Authors Contribution</i>	
Rizwan Ali Talpur	Conception of study design, acquisition, analysis, and interpretation of data.
Rabnawaz Sathio	Drafting and methodology, data interpretation
Muhammad Wasif Saleem	Analysis and interpretation of data for work

Received Date: 14-Aug-2021
 Revised Received: 22-Sept-2021
 Accepted Date: 14-Nov-2021