

Comparison of Serum level of Vitamin D in Pregnant Women with Preeclampsia and a Control Group in Isra University Hospital, Hyderabad

Nida Zaki, Uzma Parween,
Amna

ABSTRACT

Background & Objective: Conflicting results have been reported about the association between vitamin D and preeclampsia. The aim of the current study is to compare the serum levels of vitamin D in preeclamptic, and non-preeclamptic pregnant women admitted to the Isra University Hospital, Hyderabad

Materials & Methods: In this case-control study, 120 pregnant women were selected using convenience sampling from Jan 2021-2022. Then they were divided into two groups, preeclampsia and non-preeclampsia. Both groups were matched one by one. The data were collected using a demographic information questionnaire and a researcher-made checklist. After obtaining consent from participants, blood samples were collected using the peripheral venous catheter. Then the ELISA test was used to measure vitamin D in participants. Data were analyzed by SPSS version 16 using, independent t-test, and Chi-square test ($P \leq 0.05$).

Results: Of the participants, 64 (53.4 %) were between 20-35 years old, 48 (40%) were older than 35 years, and 8 (6%) were younger than 20 years. Most of the participants' gestational age was between 34 and 39 weeks (89.1%). The Mean (SD) of serum level of vitamin D was significantly higher in healthy pregnant women (19.06 (7.48)) than women with preeclampsia (16.60 (5.87)) ($P < 0.01$). Among those with preeclampsia, the highest vitamin D level (17.17) was observed at 37-39 weeks of gestation. The lowest level (10.8) was for those with less than 34 weeks gestational age.

Conclusion: This study demonstrated that vitamin D could be a protective factor for preeclampsia in pregnant women. However, further studies are recommended.

Keywords: Preeclampsia, Pregnancy, Outcomes, Vitamin D

How to cite this:

Zaki N, Parween U, Amna Comparison of Serum level of Vitamin D in Pregnant Women with Preeclampsia and a Control Group in Isra University Hospital, Hyderabad JIMC 2023; 6.(2) 403-407

Correspondence

Dr. Nida Zaki
Consultant Gynae and Obs
Isra University Hospital Hyderabad Pakistan
Email: Nidazakiqureshi@gmail.com

Introduction

Vitamin D deficiency in pregnant women is a worldwide public health problem⁽¹⁾. Vitamin D is a steroid hormone that has an important role in the innate immune system as well as several cellular activities⁽²⁾. Vitamin D deficiency during pregnancy is associated with many short- and long-term problems for fetuses and infants, including developmental delay, skeletal problems, type 1 diabetes, asthma, and schizophrenia^(3, 4).

There are two main reasons for the epidemic of vitamin D deficiency in pregnant women: insufficient sun exposure and low vitamin D intake from diet, so Vitamin D stimulates the immune response in the Bodnar *et al.* suggested vitamin D deficiency in pregnant women under 22 weeks gestation as an independent and strong risk factor for preeclampsia. This study also reported an association between reduced maternal vitamin D levels and the incidence of preeclampsia. They also reported a deficiency in neonatal serum vitamin D as a result of maternal vitamin D deficiency⁽¹⁶⁾. Many hypotheses have been suggested to justify the pathogenesis of preeclampsia. In this regard, vitamin D deficiency has been emphasized as a stimulating agent of vascular and systemic inflammation that causes insufficient placental growth^(17, 18). However, due to the complex nature of preeclampsia, there is no clear association between vitamin D deficiency and the incidence of preeclampsia. A systematic review of clinical trials investigating the impact of vitamin D supplementation has

reported a positive effect on reducing the risk of preeclampsia^(19, 20). On the other hand, some studies reported the role of vitamin D in improving endothelial function during placenta replacement. However, in vitro studies have reported the negative impact of vitamin D on the renin-angiotensin system. These studies showed a significant relationship between vitamin D and blood pressure, although the main mechanism and its exact role in hypertension have not been determined yet^(21, 22). The current study aims to compare serum levels of vitamin D in pregnant women with preeclampsia admitted to the Isra University Hospital, Hyderabad, Iran, with a control group.

decidua tissue of the placenta, which results in its healthier activity⁽⁷⁾. Some studies mentioned the imbalance of placental biomarkers and vitamin D deficiency as risk factors for preeclampsia^(8, 9). Preeclampsia is a multifactorial pregnancy complication that develops after 20 weeks of gestation^(10, 11).

The higher prevalence and more severe cases of preeclampsia in black women strengthen the association between vitamin D and preeclampsia^(12, 13). Some studies reported a seasonal pattern of preeclampsia in such a way that the lowest incidence is in summer when the serum concentration of 25(OH)D is at its highest, and its peak is in winter when the serum concentration of 25(OH)D is at its lowest^(14, 15).

Methods

In this case-control study, 240 pregnant women

admitted from October 2018 to October 2019 were divided into two groups of preeclampsia (case) and non-preeclampsia (control) (each with 120 subjects). Participants were selected using the convenience sampling technique. Participants of both groups were matched one by one. Inclusion criteria were pregnant women aged 14-45 years from the beginning of 21 weeks of gestational to the end of 40 weeks (based on nuchal translucency scan, or 12-scan, and LMP criteria for patients without ultrasound), living in the Sindh province, preeclampsia for those in the case group, and pregnancy without preeclampsia for those in the control group. The exclusion criteria included unwillingness to participate and having kidney disease, malabsorption syndromes, Vitamin D toxicity, Metastatic cancers, hypo and hyper thyroids, lupus, rheumatoid arthritis, and natural pregnancy (no IVF). Data were collected using a demographic information questionnaire and a researcher-made checklist, which included gestational age, number of pregnancies, history of chronic diseases, history of preeclampsia, and history of preeclampsia in the family members. The validity of the checklist was assessed by ten faculty

Results

The youngest and oldest participants were 14 and 45 years old, respectively. The mean serum level of vitamin D was not significantly associated with age, gestational week, and history of preeclampsia in previous pregnancies (Table 1).

In general comparison, the mean vitamin D level of those in the case group (16.60 ± 5.87 ng/mL) was lower than the control group (19.06 ± 7.48 ng/mL), which was statistically significant ($P < 0.05$).

According to the findings, out of 240 participants, 23 (9.58%) had a severe deficiency, 174 (72.5%) mild deficiency, and 43 (35.84%) had an appropriate vitamin D level. A significant difference between the two groups in terms of serum level of vitamin D was observed. Those in the preeclampsia group, had significantly lower serum level of vitamin D ($P = 0.01$). The majority of participants in the case (80.84%) and control (64.16%) groups had mild vitamin D deficiency.

Discussion

According to the results, the level of vitamin D in pregnant women serum with preeclampsia was significantly lower than non-preeclampsia pregnant women. It was not significantly associated with age, gestational week, and history of preeclampsia.

According to the Hamedanian *et al.* study conducted in Karaj, Iran, the serum level of vitamin D in pregnant women with preeclampsia was lower than in healthy pregnant women. They found no correlation between vitamin D deficiency and predicting factors of preeclampsia⁽²³⁾. A cohort study conducted in Pennsylvania, with a sample size of 2327 subjects, reported that increased serum level of vitamin D up to 50 ng/L was associated with decreased risk of preeclampsia, and in amounts above 50 ng, this risk remained constant (test of nonlinearity $P < 0.05$)⁽²⁴⁾. Achkar *et al.*, in a study conducted in Canada, also reported that preeclampsia was significantly higher in pregnant women with serum levels less than 30 nmol/L compared with those with at least 50 nmol/L of vitamin D

members of the obstetrics and gynecology department of the Isra University Hospital, Hyderabad. The goals of the study were explained to all participants, and then we obtained written consent from them. To avoid additional interventions, blood samples were collected using the patient's peripheral venous catheter. The collected samples were stored in the refrigerator and transferred to the laboratory within 20 minutes. 24 ng/L), and appropriate (24.1 to 80 ng/L). All participants were selected from one province to control the confounding variable of sunlight exposure. Data were analyzed using descriptive statistics. Then, samples were placed inside the incubator at 25-28°C. Finally, serum levels of vitamin D were measured by the ELISA test. The measured values were classified into three groups: severe deficiency (less than 10 ng/L), mild deficiency (10 to independent t-test, and Chi-square in SPSS version 16 (SPSS Inc., Chicago, Ill., USA). To test the normal distribution of data, the Kolmogorov-Smirnov test was applied. The significant level in this study was considered below 0.05 ($P < 0.05$).

(25). There are other studies that reported an association between low serum levels of vitamin D and preeclampsia^(26, 27).

A systematic review of clinical trials on the association of vitamin D supplementation on preeclamptic women reported that vitamin D supplementation could reduce the risk of preeclampsia by up to 53% compared to placebo⁽²⁸⁾. In a case-control study, Bodnar *et al.* have compared two groups of preeclampsia and control. They have investigated the serum level of D(OH)22 during pregnancy and before the onset of preeclampsia and women with preeclampsia. They reported that, in women with preeclampsia, at the initial weeks of pregnancy, the serum level of D(OH)22 was lower than the control group. These authors reported that when vitamin D levels reduced to 22 ng/L, the risk of preeclampsia increased by two folds. According to their report, maternal vitamin D deficiency could be an independent risk factor of preeclampsia⁽¹⁾.

However, Burris *et al.*, in a study performed in the United States, found no statistically significant association between preeclampsia and vitamin D levels. Also, in contrast to the hypothesis of this study, increasing maternal vitamin D levels were associated with an increased chance of developing preeclampsia. At the end of their paper, Burris *et al.* mentioned that this finding is most likely due to chance and does not indicate a causal association between vitamin D and preeclampsia. They recommended performing clinical trials to investigate this issue⁽²⁹⁾ further. It seems that vitamin D can prevent preeclampsia. Three justifications for this issue are as follows: adequate intake of vitamin D can stabilize calcium homeostasis and consequently reduce blood pressure⁽³⁰⁾ or directly suppress the smooth muscles of the vessels⁽³¹⁾. Second, vitamin D can act as an endocrine inhibitor of the renin-angiotensin system, which has a major role in regulating blood pressure. Third, as a balancer of T helper cell activity, it plays a role in strengthening the immune system⁽³²⁾.

Table 1. Mean serum level of vitamin D of participants separated by gestational age (less than 37 weeks and above 37weeks).

	to 41 weeks			
Mother age	Younger than 20 years	8	12/42	3/43
	20 to 35 years	64	18/28	5/61
	Older than 35 years	48	15/06	5/83
History of preeclampsia	Yes	99	16/77	6/13
	No	21	1578	4/44

Table 2. Comparison of vitamin D levels in case and control groups based on independent t-test

Group	Vitamin D (ng/mL)	Frequency	Percentage	P-value*
Case	Severe deficiency (< 10 ng/L)	11	9/16	
	Mild deficiency (10 to 24 10 ng/L)	97	80/84	
	Appropriate (1/24 to 10 ng/L 80)	12	10/0	2/83
Control	Severe deficiency (< 10 ng/L)	12	10/00	0/001
	Mild deficiency (10 to 24 10 ng/L)	77	64/16	
	Appropriate (1/24 to 10 ng/L 80)	31	25/84	
Variable		Frequency	Mean vitamin D (ng/ml)	Standard deviation
Gestational week	Less than 34	1	10/8	-
	34 to 37	54	16/73	5/32
	37 weeks and 1 day Up to 39 weeks	53	17/17	6/10

In the present study, there was no significant association between gestational age and serum level of vitamin D. However, according to a study that has compared the risk of preeclampsia in 12-18 and 24-26 weeks of pregnancy, in cases with vitamin D deficiency, low levels of D(OH)25 in the late second trimester was associated with an increased risk of preeclampsia. So that there was a remarkable dose-response relationship between serum levels of D(OH)25 of maternal plasma during 24-26 weeks of gestation with the risk of preeclampsia ($P=0.03$). Although this association was not significant at the gestational age of 12-18 weeks ($P=0.16$) Wei *et al.* reported that in pregnant women, serum levels of D(OH) lower than 50 nmol/L at 24-26 weeks are associated with a 3.2-fold increase in the risk of preeclampsia (OR=24.3; CI 95% 1.37-7.69). This association was not significant at gestational weeks of 12-18 (OR=24.1; CI 95% 0.58-2.67). Also, there are

other case-control studies that reported no significant association between the serum levels of mothers and the risk of developing preeclampsia during the first trimester of pregnancy⁽³³⁻³⁶⁾.

Conclusion

According to the results of the present study and other research, there is a significant association between the serum level of vitamin D during pregnancy and preeclampsia. Considering the importance of this issue and the ease of treatment and prevention of vitamin deficiency in pregnant women, it is recommended to perform clinical trials with special emphasis on gestational age and other confounding variables.

Conflict of interest: None.

Financial Support: None.

Reference

- Bodnar LM, Simhan HN, Powers RW, Frank MP, Cooperstein E, Roberts JM. High prevalence of vitamin D insufficiency in black and white pregnant women residing in the northern United States and their neonates. *J Nutr.* 2007;137(2):447-52 [DOI:10.1093/jn/137.2.447] [PMID] [PMCID]
- Chirumbolo S, Björklund G, Sboarina A, Vella A. The Role of Vitamin D in the Immune System as a Pro-survival Molecule. *Clin Ther.* 2017;39(5):894-916. [DOI:10.1016/j.clinthera.2017.03.021] [PMID]
- Yu CK, Sykes L, Sethi M, Teoh TG, Robinson S. Vitamin D deficiency and supplementation during pregnancy. *Clin Endocrinol (Oxf).* 2009;70(5):685-90. [DOI:10.1111/j.1365-2265.2008.03403.x] [PMID]
- Barrera D, Avila E, Hernandez G, Halhali A, Biruete B, Larrea F, et al. Estradiol and progesterone synthesis in human placenta is stimulated by calcitriol. *J Steroid Biochem Mol Biol.* 2007;103(3-5):529-32. [DOI:10.1016/j.jsbmb.2006.12.097] [PMID]
- Lee JM, Smith JR, Philipp BL, Chen TC, Mathieu J, Holick MF. Vitamin D deficiency in a healthy group of mothers and newborn infants. *Clin Pediatr.* 2007;46(1):42-4. [DOI:10.1177/0009922806289311] [PMID]
- Barger-Lux MJ, Heaney RP, Dowell S, Chen TC, Holick MF. Vitamin D and its major metabolites serum levels after graded oral dosing in healthy women. *Osteoporos Int.* 1998;8(3):222-30. [DOI:10.1007/s001980050058] [PMID]
- from normal and preeclamptic pregnancies. *Am J Physiol-Endocrinol Metab.* 2012;303(7):E928-35. [PMCID] [DOI:10.1152/ajpendo.00279.2012] [PMID]
- Tamblyn JA, Hewison M, Wagner CL, Bulmer JN, Kilby MD. Immunological role of vitamin D at the maternal-fetal interface. *J Endocrinol.* 2015;224(3):R107-21. [DOI:10.1530/JOE-14-0642] [PMID]
- Shin JS, Choi MY, Longtine MS, Nelson DM. Vitamin D effects on pregnancy and the placenta. *Placenta.* 2010;31(12):1027-34. [PMCID] [DOI:10.1016/j.placenta.2010.08.015] [PMID]
- Alvarez-Fernandez I, Prieto B, Rodriguez V, Ruano Y, Escudero AI, Alvarez FV. Role of vitamin D and sFlt-1/PlGF ratio in the development of early- and late-onset preeclampsia. *Clin Chem Lab Med.* 2015;53(7):1033-40. [DOI:10.1515/cclm-2014-1039] [PMID]
- Shah A, Fawole B, M'umunya JM, Amokrane F, Nafiu I, Wolomy JJ, et al. Cesarean delivery outcomes from the WHO global survey on maternal and perinatal health in Africa. *Int J Gynaecol Obstet.* 2009;107(3):191-7. [DOI:10.1016/j.ijgo.2009.08.013] [PMID]
- McClure EM, Saleem S, Pasha O, Goldenberg RL. Stillbirth in developing countries: a review of causes, risk factors and prevention strategies. *J Matern. Fetal Neonatal Med.* 2009;22(3):183-90 [DOI:10.1080/14767050802559129] [PMID] [PMCID]
- Eskenazi B, Fenster L, Sidney S. A multivariate analysis of risk factors for preeclampsia. *JAMA.* 1991;266(2):237-41. [DOI:10.1001/jama.266.2.237] [PMID]
- MacKay AP, Berg CJ, Atrash HK. Pregnancy-related mortality from preeclampsia and eclampsia. *Obstet Gynecol.* 2001;97(4):533-8. [DOI:10.1097/00006250-200104000-00011] [PMID]
- Bodnar LM, Catov JM, Roberts JM. Racial/ethnic differences in the monthly variation of preeclampsia incidence. *Am J Obstet Gynecol.* 2007;196(4):324-e1. [DOI:10.1016/j.ajog.2006.11.028] [PMID]
- Magnus P, Eskild A. Seasonal variation in the occurrence of pre-eclampsia. *Br J Obstet Gynaecol.* 2001;108(11):1116-9. [PMID] [DOI:10.1111/j.1471-0528.2003.00273.x]
- Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab.* 2007;92(9):3517-22. [DOI:10.1210/jc.2007-0718] [PMID] [PMCID]
- Ma R, Gu Y, Zhao S, Sun J, Groomer LJ, Wang Y. Expressions of vitamin D metabolic components VDBP, CYP2R1, CYP27B1, CYP24A1, and VDR in placentas
- Brodowski L, Burlakov J, Myerski AC, von Kaisenberg CS, Grundmann M, Hubel CA, et al. Vitamin D Prevents Endothelial Progenitor Cell Dysfunction Induced by Sera from Women with Preeclampsia or Conditioned Media from Hypoxic Placenta. *PLoS ONE.* 2014;9(6):e98527. [DOI:10.1371/journal.pone.0098527] [PMID] [PMCID]
- Khaing W, Vallibhakara SA, Tantrakul V, Vallibhakara O, Rattanasiri S, McEvoy M, et al. Calcium and vitamin D supplementation for prevention of preeclampsia: a systematic review and network meta-analysis. *Nutrients.* 2017;9(10):1141. [DOI:10.3390/nu9101141] [PMID] [PMCID]
- Anothaisintawee T, Attia J, Nickel JC, Thammakraisorn S, Numthavaj P, McEvoy M, et al. Management of chronic prostatitis/chronic pelvic pain syndrome: a systematic review and network meta-analysis. *JAMA.* 2011;305(1):78-86. [DOI:10.1001/jama.2010.1913] [PMID]
- Ganguly A, Tamblyn JA, Finn-Sell S, Chan SY, Westwood M, Gupta J, et al. Vitamin D, the placenta and early pregnancy: effects on trophoblast function. *J Endocrinol.* 2018;236(2):R93-103. [DOI:10.1530/JOE-17-0491] [PMID]
- Ma Y, Johnson CS, Trump DL. Mechanistic insights of vitamin D anticancer effects. *Vitamins Hormones.* 2016;100:395-431. [DOI:10.1016/bs.vh.2015.11.003] [PMID]
- Hamedanian L, Bادهnoosh B, Razavi-Khorasani N, Mohammadpour Z, Mozaffari-Khosravi H. Evaluation of vitamin D status, parathyroid hormone, and calcium among Iranian pregnant women with preeclampsia: A case-control study. *Int J Reprod BioMed.* 2019;17(11):831. [PMCID] [DOI:10.18502/ijrm.v17i10.5494] [PMID]
- Baca KM, Simhan HN, Platt RW, Bodnar LM. Low maternal 25-hydroxyvitamin D concentration increases the risk of severe and mild preeclampsia. *Ann Epidemiol.* 2016;26(12):853-7. [PMCID] [DOI:10.1016/j.annepidem.2016.09.015] [PMID]
- Achkar M, Dodds L, Giguère Y, Forest JC, Armson BA, Woolcott C, et al. Vitamin D status in early pregnancy and risk of preeclampsia. *Am J Obstet Gynecol.* 2015;212(4):511-e1. [PMCID] [DOI:10.1016/j.ajog.2014.11.009] [PMID]
- Robinson CJ, Alanis MC, Wagner CL, Hollis BW, Johnson DD. Plasma 25-hydroxyvitamin D levels in early-onset severe preeclampsia. *Am J Obstet Gynecol.* 2010;203(4):366-e1. [PMCID] [DOI:10.1016/j.ajog.2010.06.036] [PMID]
- Rezaei M, Farhadifar F, Shahgheibi S, Hojati N. Comparison of serum vitamin D levels in pregnant women with preeclampsia with healthy pregnant women. *Iran J Obstet Gynecol Infertil.* 2014;17(113):1-6. [DOI:10.22038/ijogi.2014.3384]
- Khaing W, Vallibhakara SA, Tantrakul V, Vallibhakara O, Rattanasiri S, McEvoy M, et al. Calcium and vitamin D supplementation for prevention of preeclampsia: a systematic review and network meta-analysis. *Nutrients.* 2017;9(10):1141. [DOI:10.3390/nu9101141] [PMID] [PMCID]
- Burris HH, Rifas-Shiman SL, Huh SY, Kleinman K, Litonjua AA, Oken E, et al. Vitamin D status and hypertensive disorders in pregnancy. *Ann Epidemiol.* 2014;24(5):399-403. [PMCID] [DOI:10.1016/j.annepidem.2014.02.001] [PMID]
- Villar J, Belizan JM, Fischer PJ. Epidemiologic observations on the relationship between calcium intake and eclampsia. *Int J Gynaecol Obstet.* 1983; 21(4):271-8.
- Cardus A, Parisi E, Gallego C, Aldea M, Fernandez E, Valdivielso JM. 1,25-Dihydroxyvitamin D3 stimulates vascular smooth muscle cell proliferation through a vegf-mediated
- Evans KN, Bulmer JN, Kilby MD, Hewison M. Vitamin D and placental-decidual function. *J Soc Gynecol Investig.* 2004;11(5):263-71. [DOI:10.1016/j.jsigi.2004.02.002] [PMID]
- Wei SQ, Audibert F, Hidirolou N, Sarafin K, Julien P, Wu Y, et al. Longitudinal vitamin D status in pregnancy and the risk of preeclampsia. *Int J Obstet Gynaecol.* 2012;119(7):832-9. [DOI:10.1111/j.1471-0528.2012.03307.x] [PMID]
- Powe CE, Seely EW, Rana S, Bhan I, Ecker J, Karumanchi SA, et al. First trimester vitamin D, vitamin D binding protein, and subsequent preeclampsia. Hypertension. 2010;56(4):758-63. [DOI:10.1161/HYPERTENSIONAHA.110.1582 38] [PMID] [PMCID]
- Yu CK, Ertl R, Skyfta E, Akolekar R, Nicolaidis KH. Maternal serum vitamin D levels at 11-13 weeks of gestation in preeclampsia. *J Hum Hypertens.* 2013;27(2):115-8. [DOI:10.1038/jhh.2012.1] [PMID]
- Fernández-Alonso AM, Dionis-Sánchez EC, Chedraui P, González-Salmerón MD, Pérez-López FR. First-trimester maternal serum 25-hydroxyvitamin D3 status and pregnancy outcome. *Int J Gynecol Obstet.* 2011;116(1):6-9. [DOI:10.1016/j.ijgo.2011.07.029] [PMID]

Authors

1. Dr. Nida Zaki
MS gynae and Obs
Assistant Professor
Consultant Gynae and Obs
Isra University Hospital HyderabadPakistan
2. Dr. Uzma Parveen
FCPS Gynae and obsAssociate Professor
Bilawal Medical College HyderabadPakistan
3. Dr. Amna ‘
FCPS Gynae and Obs Senior Registrar
Isra University Hospital HyderabadPakistan

Authors Contribution	
Nida Zaki	Analysis and Interpretation of data for work
Uzma Parveen	Conception of study design, acquisition, analysis, and interpretation of data.
Amna	Drafting and methodology, data interpretation

Received Date: 3-Mar-2023
Revised Received: 17-May-2023
Accepted Date: 23-Sept-2023